

Abstract for the Second Canadian International Workshop on High-Field Laser Plasma Interaction Physics, Banff, Alberta Feb 22-25 1995.

Nonlinear frequency shifts and harmonic generation in finite amplitude ion acoustic waves.

Edward A Williams,²

*Lawrence Livermore National Laboratory,
L472, Box 5508, Livermore Ca 94550*

We investigate analytically the effects of particle trapping on finite amplitude ion acoustic waves in mixed ion-species plasmas. Following Andreev and Tikhonchuk¹, we calculate the particle distribution functions in the presence of a periodic potential by solving the appropriate kinetic equation for a time long compared to the trapping time, but short compared to a collision time. The resulting distributions are flattened in the vicinity of the ion acoustic phase velocity and have trapped and untrapped components. Expressions for the perturbed charge density are obtained by integrating over velocity space and expanding in powers of $(e\phi/T)^{1/2}$. Substituting these in Poisson's equation gives an amplitude dependent dispersion relation and harmonic generation coefficients.

From this one can derive a candidate model for the saturation of SBS ion-acoustic waves.

(1) A. A. Andreev and V. T. Tikhonchuk, Sov. Phys. JETP, **68**, 1137 (1989)

(2) Work performed under the auspices of the United States Dept. of Energy by the Lawrence Livermore National Laboratory under contract number W-7405-ENG-48.